

SID 16
(3 hits this paper)

RESULT 2
BTU51820
LOCUS BTU51820 376 bp DNA MAM 16-MAY-1996
DEFINITION Bos taurus Hereford 7 mitochondrial hypervariable region D-loop,
partial sequence.
ACCESSION U51820
VERSION U51820.1 GI:1314616
KEYWORDS .
SOURCE cow strain=Hereford.
ORGANISM Mitochondrion Bos taurus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
Bovidae; Bovinae; Bos.
REFERENCE 1 (bases 1 to 376)
AUTHORS Bradley, D.G., MacHugh, D.E., Cunningham, P. and Loftus, R.T.
TITLE Mitochondrial diversity and the origins of African and European
cattle
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 93 (10), 5131-5135 (1996)
MEDLINE 96209869
REFERENCE 2 (bases 1 to 376)
AUTHORS Bradley, D.G., MacHugh, D.E., Cunningham, P. and Loftus, R.T.
TITLE Direct Submission
JOURNAL Submitted (19-MAR-1996) Daniel G. Bradley, Genetics, Trinity
College Dublin, Dublin, 2, Ireland
FEATURES Location/Qualifiers
source 1. .376
/organism="Bos taurus"
/organelle="mitochondrion"
/strain="Hereford"
/db_xref="taxon:9913"
/note="HE7"
D-loop <1. .>376
/note="hypervariable region"
BASE COUNT 106 a 92 c 60 g 118 t
ORIGIN

Query Match 100.0%; Score 153; DB 3; Length 376;
Best Local Similarity 100.0%; Pred. No. 1e-23;
Matches 153; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 gcatataagcaagtacatgacctctatagcagtacataatacatataattattgactgta 60
|||||
Db 4 GCATATAAGCAAGTACATGACCTCTATAGCAGTACATAATACATATAATTATTGACTGTA 63

Qy 61 catagtacattatgtcaaattcattcttgatagtatatctattatatattccttaccatt 120
|||||
Db 64 CATAGTACATTATGTCAAATTCATTCTTGATAGTATATCTATTATATATTCCTTACCATT 123

Qy 121 agatcacgagcttaattaccatgccgcgtgaaa 153
|||||
Db 124 AGATCACGAGCTTAATTACCATGCCGCGTGAAA 156

sequencer. Consensus sequences were constructed and analyzed with the help of the University of Wisconsin GCG software package. The BLAST algorithm from the National Center for Biotechnology Information (NCBI) was also employed for nucleotide and amino acid sequence homology search (Altschul, S. F. et al. (1990) *J. Mol. Biol.* 215:403-410).

Preparation of Recombinant Proteins:

The recombinant CR8 protein with histidine-tag was prepared using the Xpress™ System (Invitrogen) according to the manufacturer's protocol. cDNA corresponding to the CR8 bHLH domain was obtained by PCR. The sequences of the primers, 5'-GGGGTCTACCAGGGATGTAC-3' (SEQ ID NO:15) for the 5' side, and 5'-GTAAACCACTCTGCAGGGCAATGA-3' (SEQ ID NO:16) for the 3' side, were slightly different from the final consensus sequence for CR8, but the difference did not affect the core bHLH motif. The PCR product was cloned into pT7Blue T-vector (Novagen) and subsequently into pRSET-A vector at BamIII and HindIII sites. Constructs were confirmed by DNA sequencing. The protein was overexpressed in JM109 at 37° C. in the presence of isopropylthio- β -D-galactoside (IPTG) by infecting the bacteria with M13 phages that contain the T7 RNA polymerase gene. The cells were lysed with 100 μ g/ml lysozyme in native binding buffer (20 mM sodium phosphate, pH 7.8, 500 mM NaCl), the lysate was loaded on a ProBond™ Ni2+ column, and the recombinant protein was eluted with native-imidazole elution buffer (20 mM sodium phosphate, pH 6.0, 500 mM NaCl, 500 mM imidazole). The protein was then dialyzed against lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 10% glycerol, 0.1% Triton X-100, 1 mM DTT) and concentrated on a Microcon 10 (Amicon). To confirm the purity and the quantity, the protein was fractionated on a 12% SDS-polyacrylamide gel with protein standards of known concentrations and visualized by silver staining. The recombinant protein corresponding to the bHLH domain of *da* was supplied by Dr. Michael Caudy (Cornell University Medical College).

Mobility Shift Assay:

The oligonucleotide probes used for the electrophoretic mobility shift assay (EMSA) are as follows: the Class A site used was the CACCTG hexamer (CAGGTG for the opposite strand) from the T5 promoter region of the *Drosophila* AS-C (Villares, R. et al. (1987) *Cell* 50:415-424) (5'-GATCGTAGTCACGCAGGTGGGATCCCTA-3' (SEQ ID NO:17) and 5'-GATCTAGGGATCCACCTGCGTGACTAC-3' (SEQ ID NO:18) for the opposite strand), the Class B site was the CACGTG hexamer from the USF binding site in the adenovirus major late promoter (Gregor, P. D. et al. (1990) *Genes & Dev.* 4:1730-1740) (5'-GATCGGTGTAGGCCACGTGACCGGGTGT-3' (SEQ ID NO:19) and 5'-GATCACACCCGGTCACGTGGCCTACACC-3' (SEQ ID NO:20), the Class C site was the CACGCG hexamer (CGCGTG for the opposite strand) from the AS-C T5 promoter (5'-GATCGGCAGCCGGCACGCGACAGGGCC-3' (SEQ ID NO:21) and 5'-GATCGGCCCTGTGCGGTGCCGGCTGCC-3' (SEQ ID NO:22), and the N-box (CACNAG) was the double hexamer sequence from the Enhancer of split [E(spl)] m8 promoter (Klimbt, C. et al. (1989) *EMBO J.* 8:203-210) (5'-GATCACGCCACGAGCCACAAGGATG-3' (SEQ ID NO:23) and 5'-GATCCAATCCTTGTGGCTCGTGGCGT-3' (SEQ ID NO:24). One strand of the oligonucleotide was

labeled with [γ -³²P]ATP by T4 polynucleotide kinase, hybridized with three times excess of the opposite strand, and purified using MERmaid oligonucleotide purification kit (BIO 101). 150 ng of the protein was allowed to bind to 50,000 cpm (equivalent to 0.5 ng in a typical experiment) of the labeled probe for 15 min at room temperature in 20 mM Hepes, pH 7.6, 50 mM KCl, 10 mM DTT, 5% glycerol, 0.5 mM EDTA and 0.3 mg/ml BSA. Two microgram of poly (dI-dC) was added to each 20 μ l reaction as on-specific DNA. Samples were analyzed on a 5% native polyacrylamide gel and visualized by autoradiography.

Regulation of CR8:

CR genes were originally defined in IL-2 stimulated normal human T cells. To examine CR8 expression in cytokine-dependent cell lines, the level of CR8 expression was measured by Northern hybridization in the IL-2-dependent human T cell line Kit 225, the IL-3-dependent mouse pro-B cell line Ba/F3, and the IL-2-dependent mouse T cell line CTLL2. The results of this experiment are illustrated in FIGS. 13A-13C. In FIG. 13A, RNA was isolated from quiescent normal human T cells (lanes 1 and 2), IL-2-dependent human T cell line Kit 225 (lanes 3 and 4), IL-3-dependent mouse pro-B cell line Ba/F3 (lanes 5 and 6) and IL-2-dependent mouse T cell line CTLL2 (lanes 7 and 8) left untreated (lanes 1, 3, 5, and 7) or stimulated with 500 pM recombinant human IL-2 (lanes 2, 4, and 8) or 10 U/ml recombinant mouse IL-3 (lane 6) for 2 hr at 37° C. The amount of the growth factor used in the experiment was sufficient to induce maximal [³H]thymidine incorporation. Ten microgram of total RNA was analyzed on formaldehyde/agarose gel and hybridized with either human (lanes 1 to 4) or mouse (lane 5 to 8) CR8 cDNA.

As shown in FIG. 13A, a single 3.2 kb species hybridized to the cDNA probe, and in all three cell lines tested, the level of CR8 was clearly augmented when the cells were stimulated with their respective growth factors. Correlation between the level of CR8 and that of DNA synthesis was in the presence of growth-inhibitory agents was also examined. In this regard, increases in cytoplasmic cAMP are known to inhibit the growth of many cell types, including lymphocytes (Johnson, K. W. et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:6072-6076). IFNs also exert antiproliferative activity on many cell types (Pestka, S. et al. (1987) *Annu. Rev. Biochem.* 56:727-777). Therefore, Kit 225 was stimulated with IL-2, IFN β , or forskolin, which increases cytoplasmic cAMP by activating adenylate cyclase, either alone or in combination. IL-2-dependent [³H]thymidine incorporation was inhibited by IFN β and forskolin in Kit 225 cells in a dose-dependent fashion (FIG. 13B). FIG. 13B demonstrates that IFN β and forskolin inhibit IL-2-dependent [³H]thymidine incorporation by Kit 225 cells. Ten thousand quiescent Kit 225 cells were incubated with indicated reagents in 200 μ l for 24 hr at 37° C. The culture was pulsed with [³H]thymidine for the last 4 hr to monitor the DNA synthesis. (●), IL-2 only (500 pM); (◇), IL-2 500 pM+varying concentrations of IFN β (U/ml); (○), IL-2 500 pM+varying concentrations of forskolin (fM). While forskolin was capable of reducing the IL-2-dependent [³H]thymidine incorporation almost to the basal level, IFN β -mediated inhibition never exceeded 70% of the maximal incorporation in several independent experiments. The expression of CR8 was compared with that of c-myc, an IL-2-inducible immediate-early gene that encodes a bHLH protein and is implicated for cell proliferation (Marcu, K. B. et al. (1992) *Annu. Rev. Biochem.* 61:809-860).

FIG. 13C shows the effect of antiproliferative agents on the expression of CR8 and c-myc transcripts in Kit 225.

5104

RESULT 7
 BTU51824
 LOCUS BTU51824 389 bp DNA MAM 16-MAY-1996
 DEFINITION Bos taurus Jersey 5 mitochondrial hypervariable region D-loop, partial sequence.
 ACCESSION U51824
 VERSION U51824.1 GI:1314620
 KEYWORDS .
 SOURCE cow strain=Jersey.
 ORGANISM Mitochondrion Bos taurus
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea; Bovidae; Bovinae; Bos.
 REFERENCE 1 (bases 1 to 389)
 AUTHORS Bradley,D.G., MacHugh,D.E., Cunningham,P. and Loftus,R.T.
 TITLE Mitochondrial diversity and the origins of African and European cattle
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 93 (10), 5131-5135 (1996)
 MEDLINE 96209869
 REFERENCE 2 (bases 1 to 389)
 AUTHORS Bradley,D.G., MacHugh,D.E., Cunningham,P. and Loftus,R.T.
 TITLE Direct Submission
 JOURNAL Submitted (19-MAR-1996) Daniel G. Bradley, Genetics, Trinity College Dublin, Dublin, 2, Ireland
 FEATURES Location/Qualifiers
 source 1. .389
 /organism="Bos taurus"
 /organelle="mitochondrion"
 /strain="Jersey"
 /db_xref="taxon:9913"
 /note="JE5"
 D-loop <1. .>389
 /note="hypervariable region"
 BASE COUNT 111 a 93 c 61 g 124 t
 ORIGIN

Query Match 100.0%; Score 46; DB 3; Length 389;
 Best Local Similarity 100.0%; Pred. No. 1.8e-08;
 Matches 46; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 attatatgccccatgcatataagcaagtacatgacctctatagcag 46
 ||||||||||||||||||||||||||||||||||||||||||||
 Db 3 ATTATATGCCCCATGCATATAAGCAAGTACATGACCTCTATAGCAG 48

2 111 from 1996 paper

-continued

AAGCCAGCCG GGAAGAGAGG TTGAGTCACC CATCCCCGAG GTGGCTGCCC CTGTGTGGGA	802
GGCAGGTTCT GCAAAGCAAG TGCAAGAGGA CAAAAA AAAA AAAAATGCGC	862
TCCAGCAGCC TGTTTGGGAA GCAGCAGTCT CTCCTTCAGA TACTGTGGGA CTCATGCTGG	922
AGAGGAGCCG CCCACTTCCA GGACCTGTGA ATAAGGGCTA ATGATGAGGG TTGGTGGGGC	982
TCTCTGTGGG GCAAAAAGGT GGTATGGGGG TTAGCACTGG CTCTCGTTCT CACCGAGAA	1042
GGAAGTGTC TAGTGTGGTT TAGGAAACAT GTGGATAAAG GGAACCATGA AAATGAGAGG	1102
AGGAAAGACA TCCAGATCAG CTGTTTGGCC TGTGCTCAG TTGACTCTGA TTGCATCCTG	1162
TTTTCTANT TCCCAGACTG TTCTGGGCAC GGAAGGSACC CTGGATGTGG AGTCTTCCCC	1222
TTTGGCCCTC CTCAGTGCC TCTGGGCTAG CCCAGAGTCC CTTAGCTTGT ACCTCGTAAC	1282
ACTCCTGTGT GTCTGTCCAG CCTTGCAGTC ATGTCAAGGC CAGCAAGCTG ATGTGACTCT	1342
TGCCCCATGC GAGATATTTA TACCTCAAAC ACTGGCCTGT GAGCCCTTC CAAGTCAGTG	1402
GAGAGCCCTG AAAGGAGCCT CACTTGAATC CAGCTCAGTG CTCTGGGTGG CCCCTGCAG	1462
GTGCCCCCTG ACCCTGCGTT GCAGCAGGGT CCACCTGTGA GCAGGCCCGC CCTGGGCCCT	1522
CTTCTGGAT GTGCCCTCTC TGAGTTCTGT GCTGTCTCTT GGAGGCAGGG CCCAGGAGAA	1582
CAAAGTGTGG AGGCCTCGGG GAGTGACTTT TCCAGCTCTC ATGCCCCGCA GTGTGAACA	1642
AGGCAGAAAA GGATCCTAGG AAATAAGTCT CTTGGCGGTC CCTCAGAGTC CTGCTGAAAT	1702
CCAGCCAGTG TTTTTTGTGG TATGAGAACA GCCAAAAGA GATGCCCGA GATAGAAGGG	1762
GAGCCTGTGT TTTCTTCTC GCAGACGTGA GATGAACACT GGAGTGGGCA GAGGTGGCCC	1822
AGGACCATGA CACCCTTAGA GTGCAGAAGC TGGGGGAGA GGCTGCTTCG AAGGGCAGGA	1882
CTGGGGATAA TCAGAACCTG CCTGTACCT CAGGGCATCA CTGAACAAAC ATTTCTGAT	1942
GGGAACCTCT GCGGCAGAGC CCAGGCTGGG GAAGTGAAC ACCCAGGCA GCCCCTTTGT	2002
GGCCAGGAT AATCAACACT GTTCTCTCTG TACCATGAGC TCCTCCAGGA GATTATTAA	2062
GTGTATTGTA TCATTGGTTT TCTGTGATTG TCATAACATT GTTTTGTTA CTGTTGGTGC	2122
TGTTGTTAT TATTATTGTA ATTCAGTTT GCCTCTACTG GAGAATCTCA GCAGGGGTTT	2182
CAGCCTGACT GTCTCCCTTT CTCTACCAGA CTCTACCTCT GAATGTGCTG GGAACCTCTT	2242
GGAGCCTGTC AGGAACCTCT CACTGTTTAA ATATTAGGT ATTGTGACAA ATGGAGCTGG	2302
TTTCTAGAA ATGAATGATG TTTGCAATCC CCATTTTCCT GTTTCAGCAT GTTATATTCT	2362
TATGAAATAA AAGCCCAAGT CCAATATGAA AAAAAAAAAA AAAA	2406

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 202 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Cys	Arg	Thr	Leu	Ala	Ala	Phe	Pro	Thr	Cys	Leu	Glu	Arg	Ala	
1				5					10				15		
Lys	Glu	Phe	Lys	Thr	Arg	Leu	Gly	Ile	Phe	Leu	His	Lys	Ser	Glu	Leu
			20					25					30		
Gly	Cys	Asp	Thr	Gly	Ser	Thr	Gly	Lys	Phe	Glu	Trp	Gly	Ser	Lys	His
			35					40					45		
Ser	Lys	Glu	Asn	Arg	Asn	Phe	Ser	Glu	Asp	Val	Leu	Gly	Trp	Arg	Glu
			50				55						60		